

VELAMENTOUS CORD INSERTION: SIGNIFICANCE OF PRENATAL DETECTION TO PREDICT PERINATAL COMPLICATIONS

Junichi Hasegawa*, Ryu Matsuoka, Kiyotake Ichizuka, Akihiko Sekizawa, Takashi Okai

Department of Obstetrics and Gynecology, Showa University School of Medicine, Tokyo, Japan.

SUMMARY

In the maternal and child health statistics of Japan for 2003, perinatal deaths were most frequent in pregnant women with abnormalities of the placenta, umbilical cord, and fetal membrane. Despite advances in perinatal medicine, approximately 2% of low-risk pregnant women still require an emergency cesarean section after the onset of labor. Because it is likely that half of these cases are associated with placental and umbilical cord abnormalities, it is thought that prenatal detection of such abnormalities would reduce the number of emergency cesarean sections in low-risk women. In our previous studies, some abnormalities of the placenta and umbilical cord were associated with abnormalities of cord insertion. Furthermore, we reported that prenatal detection of velamentous cord insertion (VCI) reduced the number of emergency cesarean sections in low-risk women. In this review, we describe the prenatal detection of abnormalities of umbilical cord insertion and the management of VCI based on our current clinical data. [*Taiwanese J Obstet Gynecol* 2006;45(1): 21-25]

Key Words: velamentous cord insertion, vasa previa, umbilical cord, prenatal diagnosis, ultrasound, placenta previa, variable deceleration

Velamentous Cord Insertion

The umbilical cord normally inserts into the central portion of the placenta, well away from the placental edge. The site of placental cord insertion (CI) is considered abnormal when it is located at the edge of the placental disk (marginal CI) or when the umbilical vessels separate from each other and course between the amnion and chorion before reaching the placenta (velamentous CI, VCI). The rate of VCI ranges from 0.5% to 1.69% of singleton pregnancies [1,2]. It has been reported that abnormal CI is associated with fetal growth restriction, preterm labor, abnormal intrapartum

fetal heart rate (FHR) pattern, low Apgar scores at 1 and 5 minutes, neonatal death [1,2], and abruptio placenta [3-5].

In cases with VCI, it is thought that the abnormal FHR pattern and some perinatal complications are caused by lack of Wharton's jelly, which results in compression of vessels during uterine contraction or fetal movement. When the CI site is in the lower uterine segment, VCI and marginal CI are strongly associated with variable decelerations, non-reassuring fetal status, emergency cesarean sections, and other perinatal complications. In our previous report, variable decelerations were observed in 19 of 30 (63%) upper-middle VCIs and four of five (80%) lower VCIs [6,7]. A non-reassuring FHR pattern was observed in four (13.3%) upper-middle VCIs and four (80%) lower VCIs. Emergency cesarean sections were performed in three (10%) upper-middle VCIs and three (60%) lower VCIs. Furthermore, aberrant vessels in lower VCI cases tend to become longer than those in middle and upper VCI (the length of the aberrant vessels was 3.9 ± 3.3 cm in upper-segment

*Correspondence to: Dr. Junichi Hasegawa, Department of Obstetrics and Gynecology, Showa University School of Medicine, 1-5-8 Hatanodai, Shinagawa-ku, Tokyo, 142-8666, Japan.

E-mail: hasejun@oak.dti.ne.jp

Received: February 6, 2006

Revised: February 13, 2006

Accepted: February 14, 2006

VCI, 4.7 ± 4.6 cm in middle-segment VCIs, and 10.6 ± 6.8 cm in lower-segment VCIs), because aberrant vessels are elongated by extension of the uterine isthmus as gestation advances [6,7]. It appears that the lower and longer aberrant vessels of VCI are readily compressed by the fetus, which results in an abnormal FHR pattern and intrapartum complications during labor.

In our previous study, two of 35 VCI cases (6%) and one of 39 marginal CI cases (3%) had placental abruption [6], although the incidence is 0.6–1.0% of all pregnancies [3–5]. Toivonen et al also reported that VCI has a significantly higher risk for placental abruption with an odds ratio of 2.53 [5]. Lower VCI may be associated with extension of the lower uterine segment and atrophy of the chorion villosus that covers the lower segment of the uterus, resulting in abruptio placenta, placenta accreta, accessory placenta, or placental infarction.

In multiple pregnancies, the prevalence of VCI is 10-fold higher than in singleton pregnancies. Only 56.1% of dichorionic diamniotic twins had normal CIs in both placentas. The finding was even more profound for monochorionic diamniotic (MD) twins, of which only 23.8% demonstrated normal CI [8]. Abnormalities of the umbilical cord, particularly VCI, are associated with preterm delivery, twin-to-twin transfusion syndrome (TTTS), and fetal growth restriction [1]. Hanley et al reported a 13-fold increase in the risk of birth weight discordancy in MD twins in the presence of VCI, with a rate of 46% [8]. Fries et al reported that in 38 pairs of MD twins, 11 cases (28.9%) developed TTTS. In these pregnancies, 63.6% had a VCI. The incidences of TTTS in the groups with normal CI and VCI were 18.5% and 58.3%, respectively [9]. Thus, since the influence of VCI is stronger in multiple pregnancies than in single pregnancies, it is thought that prenatal detection of VCI is more important in the management of multiple pregnancies.

Vasa Previa

Vasa previa is a condition in which the umbilical and fetal vessels are unsupported by the placenta or Wharton's jelly and traverse the region of the internal os. Vasa previa is a form of VCI in which the velamentous vessels traverse the fetal membranes in the lower uterine segment. The incidence of vasa previa is estimated at 1:50 VCI cases [10,11] or 1:1,200–5,000 pregnancies [12–15]. Because vasa previa has unsupported fetal vessels below the fetal presenting part, these fetal vessels are easily compressed or ruptured when uterine contractions or membrane rupture occurs, resulting in

fetal exsanguination. Since bleeding from vasa previa is of fetal origin, the associated fetal morbidity and mortality are extremely high, ranging from 50% to 60% with intact membranes, and 70% to 100% with ruptured membranes [12,16,17]. An intrapartum clinical diagnosis is rarely made and the diagnosis is very difficult [18].

Catanzarite et al classified vasa previa into two types: Type I is associated with VCI and Type II is associated with vessels that connect between placental lobes and traverse the internal os. The pathophysiology of Type II vasa previa may be explained by atrophy of the partial placenta around the cervical os due to a poor blood supply [18]. They found that of their 10 cases and 23 other reported cases, 54.5% (18/33) were Type II [14,18–23].

Oyelese et al performed a multicenter study of 155 pregnancies complicated by vasa previa [24]. Between 1991 and 2003, 61 of these cases were prenatally diagnosed by ultrasonography/color Doppler and the infant survival rate was 97% (59/61). In cases that were not prenatally diagnosed, the survival rate was 44% (41/94). Multivariable logistic regression analysis revealed that the only significant predictors of neonatal survival were prenatal diagnosis and gestational age at delivery. Thus, high fetal mortality due to vasa previa can be reduced by antenatal diagnosis and elective cesarean section [14,18,24,25].

Several authors have reported vasa previa as well as other placental abnormalities associated with *in vitro* fertilization (IVF) [24,26,27]. Marginal CI and VCI were more frequently found in the IVF population than the general obstetric population. Schachter et al reported that the frequencies of vasa previa and VCI after IVF pregnancy were 1:293 and 1:167, respectively, with odds ratios compared with non-IVF pregnancy calculated as 27.4 (95% confidence interval, CI, 7.4–91.7) and 5.9 (95% CI, 2.5–13.2), respectively. Abnormal placentation and situations that may lead to abnormal placentation are risk factors for vasa previa [27]. Placenta previa and bi-lobed and succenturiate-lobed placentas are risk factors for vasa previa [13,19,23,28], and these are more common in pregnancies achieved after IVF.

Ultrasound Screening

Since pregnancies complicated by VCI, including vasa previa, are at greater risk for adverse perinatal outcome, previous investigators have suggested that systematic ultrasound evaluation of the placental CI site is extremely important [14,25,29,30]. Pretorius et al reported that the detection rate of CI was markedly influenced by




Figure 1. Ultrasound image of velamentous cord insertion in the mid-trimester.

gestational age, ranging from 67% at 15–20 weeks to 30% at 36–40 weeks [30]. As visualization of the placental CI site becomes more difficult with advancing gestation, it should be evaluated in the mid trimester [14,25,31] (Figure 1). One fetal-medicine specialist identified the placental CI site in more than 99% of cases using routine color Doppler scanning in the mid trimester [14,25]. However, in our previous study, antenatal detection of VCI at 18 weeks had a sensitivity of 63% (25/40), specificity of 100% (3406/3406), positive predictive value (PPV) of 100% (25/25), and negative predictive value (NPV) of 99.6% (3406/3421) [6]. Because a number of generalist obstetricians/gynecologists with 3–5 years' experience performed the ultrasound scans in our department, and color Doppler scanning was not used in all cases, the sensitivity of detecting VCI was not high at 18–20 weeks of gestation. Our criteria for ultrasound diagnosis of a VCI were: umbilical vessels entering the placenta margin parallel to the uterine wall and connecting to superficial placental vessels; an immobile CI, even when the uterus is shaken; and umbilical vessels diverging as they traverse the membrane. In fact, the CI site could not be determined more frequently in cases of marginal CI and VCI than in the normal CI. An abnormal CI was present more frequently when it was difficult to image the CI (decreased sensitivity); thus, more precise scanning (in different body positions and using color Doppler) is indicated.

Previous studies have reported that the risk factors for vasa previa include: ultrasound diagnosis of low-lying placenta or placenta previa at earlier gestation; bi-lobed or succenturiate placenta [13,19,23,28]; multiple gestation; suspicion of aberrant vessels [23]; VCI [23,30]; and an IVF pregnancy [24]. Of patients with vasa previa at delivery, 60–69% were diagnosed in the mid-trimester [24,28], whereas, in cases without

vasa previa, two of 52 patients (3.8%) were diagnosed as having placenta previa in the mid-trimester [28]. Thus, routine transvaginal color Doppler sonography of the cervical region during the third trimester for women with increased risk is recommended [24,28].

An abnormal CI located in the lower uterine segment is strongly associated with perinatal complications and the pregnancy should be deemed high risk. We think the assessment of CI at first-trimester sonography may improve the sensitivity of detecting CI in the lower uterine segment and have profound implications for management. We reported that in cases in which CI was noted to be in the lower uterine segment during the first trimester, regardless of the location of the placenta at delivery, developmental abnormalities of the placenta and umbilical cord, including low-lying placentas, infarction and abruption of placentas, VCI and marginal CI, occurred frequently [32] (Figure 2). Because of a high visualization rate and short time for detection of cord insertion at 9–11 weeks of gestation, we propose that the systematic identification of CI in the lower uterine segment during the first trimester is extremely useful for identification of high-risk pregnancies, including VCI located in the lower uterine segment and vasa previa. In cases where CI is low at the first-trimester screening, it should be re-evaluated during the third trimester using color Doppler ultrasonography. Through our management of pregnant women, we believe that a safer delivery can be provided with accurate identification of high-risk pregnancies with abnormalities of the placenta and umbilical cord.

Treatment of Pregnancy and Delivery

Variable decelerations and a non-reassuring FHR pattern are frequently observed in cases with VCI. The



Figure 2. Ultrasound image of cord insertion (arrow) into the lower uterine segment at 10 weeks of gestation. In Os = internal os.

increased frequency of variable decelerations is most likely due to the lack of Wharton's jelly in the umbilical vessels, which results in compression of these vessels during uterine contraction or fetal movement. It is thought that variable decelerations occur as a result of cord compression. Lee et al suggest that variation in typical variable deceleration is caused by different degrees of partial cord compression [33]. However, the cause of frequent variable decelerations in VCI is the compression of aberrant vessels and the obstruction of blood flow in both arteries and veins at the same time by the uterine contraction. Variable deceleration without baroreceptor-mediated acceleration (VDNA) frequently occurs in VCI cases (Figure 3). Analysis of VDNA occurrence in the second stage of labor shows that the rate of VDNA is about three times higher in VCI cases than in controls [7].

In cases of VCI and vasa previa, we should offer frequent FHR monitoring in the late third trimester, as long as the pregnancy is not complicated with preterm labor, bleeding, shortening of the cervix, or spontaneous rupture of membranes. Patients should be educated regarding signs and symptoms of preterm of labor. Oyelese et al recommend that women with prenatally diagnosed vasa previa should be offered elective delivery by cesarean section at about 35 weeks of gestation,

or earlier if fetal lung maturity is documented [24]. Because we consider that lower VCI is analogous to vasa previa and has a very high rate of non-reassuring fetal status, we recommend elective cesarean section in cases with lower VCI. Other VCI cases can be offered an attempt at vaginal delivery with the backup of an emergency cesarean. During labor, variable deceleration, especially VDNA, should be considered a warning sign of abrupt FHR abnormalities.

Conclusion

VCI is associated with increased rates of abnormal FHR tracings and cesarean deliveries. In particular, pregnancies with a VCI located in the lower uterine segment or vasa previa should be deemed high risk. Antenatal diagnosis of a VCI in early gestation improves obstetric management. In cases in which CI is noted to be in the lower uterine segment during the first trimester, developmental abnormalities of the placenta and the umbilical cord (including VCI and vasa previa) occur frequently. Because of a high visualization rate and short time for detection of CI, systematic identification of CI during the first trimester is extremely useful for the identification of high-risk pregnancies.




Figure 3. Fetal heart rate pattern named variable deceleration with no acceleration (VDNA).

References

1. Eddleman KA, Lockwood CJ, Berkowitz GS, Lapinski RH, Berkowitz RL. Clinical significance and sonographic diagnosis of velamentous umbilical cord insertion. *Am J Perinatol* 1992; 9:123-6.
2. Heinonen S, Ryynanen M, Kirkinen P, Saarikoski S. Perinatal diagnostic evaluation of velamentous umbilical cord insertion: clinical, Doppler, and ultrasonic findings. *Obstet Gynecol* 1996;87:112-7.
3. Ananth CV, Savitz DA, Williams MA. Placental abruption and its association with hypertension and prolonged rupture of membranes: a methodologic review and meta-analysis. *Obstet Gynecol* 1996;88:309-18.
4. Ananth CV, Smulian JC, Vintzileos AM. Incidence of placental abruption in relation to cigarette smoking and hypertensive disorders during pregnancy: a meta-analysis of observational studies. *Obstet Gynecol* 1999;93:622-8.
5. Toivonen S, Heinonen S, Anttila M, Kosma VM, Saarikoski S. Reproductive risk factors, Doppler findings, and outcome of affected births in placental abruption: a population-based analysis. *Am J Perinatol* 2002;19:451-60.
6. Hasegawa J, Matsuoka R, Ichizuka K, Sekizawa A, Farina A, Okai T. Velamentous cord insertion into the lower third of the uterus is associated with intrapartum fetal heart rate abnormalities. *Ultrasound Obstet Gynecol* 2006 (in press).
7. Hasegawa J, Matsuoka R, Ichizuka K, Sekizawa A, Farina A, Okai T. Velamentous cord insertion and atypical variable de-

- celerations with no accelerations. *Int J Gynaecol Obstet* 2005; 90:26-30.
8. Hanley ML, Ananth CV, Shen-Schwarz S, Smulian JC, Lai YL, Vintzileos AM. Placental cord insertion and birth weight discordancy in twin gestations. *Obstet Gynecol* 2002;99: 477-82.
 9. Fries MH, Goldstein RB, Kilpatrick SJ, Golbus MS, Callen PW, Filly RA. The role of velamentous cord insertion in the etiology of twin-twin transfusion syndrome. *Obstet Gynecol* 1993;81:569-74.
 10. Vago T, Caspi E. Antepartum bleeding due to injury of velamentous placental vessels. *Obstet Gynecol* 1962;20:671-4.
 11. Quek SP, Tan KL. Vasa praevia. *Aust NZ J Obstet Gynaecol* 1972;12:206-9.
 12. Pent D. Vasa previa. *Trans Pac Coast Obstet Gynecol Soc* 1979; 46:50-4.
 13. Fung TY, Lau TK. Poor perinatal outcome associated with vasa previa: is it preventable? A report of three cases and review of the literature. *Ultrasound Obstet Gynecol* 1998;12: 430-3.
 14. Nomiya M, Toyota Y, Kawano H. Antenatal diagnosis of velamentous umbilical cord insertion and vasa previa with color Doppler imaging. *Ultrasound Obstet Gynecol* 1998;12: 426-9.
 15. Oyelese KO, Turner M, Lees C, Campbell S. Vasa previa: an avoidable obstetric tragedy. *Obstet Gynecol Surv* 1999;54: 138-45.
 16. Antoine C, Young BK, Silverman F, Greco MA, Alvarez SP. Sinusoidal fetal heart rate pattern with vasa previa in twin pregnancy. *J Reprod Med* 1982;27:295-300.
 17. Naftolin F, Mishell DR Jr. Vasa previa. Report of 3 cases. *Obstet Gynecol* 1965;26:561-5.
 18. Catanzarite V, Maida C, Thomas W, Mendoza A, Stanco L, Piacquadio KM. Prenatal sonographic diagnosis of vasa previa: ultrasound findings and obstetric outcome in ten cases. *Ultrasound Obstet Gynecol* 2001;18:109-15.
 19. Oyelese KO, Schwarzler P, Coates S, Sanusi FA, Hamid R, Campbell S. A strategy for reducing the mortality rate from vasa previa using transvaginal sonography with color Doppler. *Ultrasound Obstet Gynecol* 1998;12:434-8.
 20. Hertzberg BS, Kliever MA. Vasa previa: prenatal diagnosis by transperineal sonography with Doppler evaluation. *J Clin Ultrasound* 1998;26:405-8.
 21. Chen KH, Konchak P. Use of transvaginal color Doppler ultrasound to diagnose vasa previa. *J Am Osteopath Assoc* 1998;98:116-7.
 22. Sauerbrei EE, Davies GL. Diagnosis of vasa previa with endovaginal color Doppler and power Doppler sonography: report of two cases. *J Ultrasound Med* 1998;17:393-8.
 23. Lee W, Lee VL, Kirk JS, Sloan CT, Smith RS, Comstock CH. Vasa previa: prenatal diagnosis, natural evolution, and clinical outcome. *Obstet Gynecol* 2000;95:572-6.
 24. Oyelese Y, Catanzarite V, Prefumo F, et al. Vasa previa: the impact of prenatal diagnosis on outcomes. *Obstet Gynecol* 2004;103:937-42.
 25. Sepulveda W, Rojas I, Robert JA, Schnapp C, Alcalde JL. Prenatal detection of velamentous insertion of the umbilical cord: a prospective color Doppler ultrasound study. *Ultrasound Obstet Gynecol* 2003;21:564-9.
 26. Englert Y, Imbert MC, Van Rosendael E, et al. Morphological anomalies in the placentae of IVF pregnancies: preliminary report of a multicentric study. *Hum Reprod* 1987;2:155-7.
 27. Schachter M, Tovbin Y, Arieli S, Friedler S, Ron-El R, Sherman D. In vitro fertilization is a risk factor for vasa previa. *Fertil Steril* 2002;78:642-3.
 28. Francois K, Mayer S, Harris C, Perlow JH. Association of vasa previa at delivery with a history of second-trimester placenta previa. *J Reprod Med* 2003;48:771-4.
 29. Di Salvo DN, Benson CB, Laing FC, Brown DL, Frates MC, Doubilet PM. Sonographic evaluation of the placental cord insertion site. *AJR Am J Roentgenol* 1998;170:1295-8.
 30. Pretorius DH, Chau C, Poeltler DM, Mendoza A, Catanzarite VA, Hollenbach KA. Placental cord insertion visualization with prenatal ultrasonography. *J Ultrasound Med* 1996;15: 585-93.
 31. Oyelese Y. Placenta previa and vasa previa: time to leave the Dark Ages. *Ultrasound Obstet Gynecol* 2001;18:96-9.
 32. Hasegawa J, Matsuoka R, Ichizuka K, Otsuki K, Sekizawa A, Farina A, Okai T. Cord insertion into the lower uterine segment in first-trimester is associated with placenta and umbilical cord abnormalities. *Ultrasound Obstet Gynecol* (In press).
 33. Lee CY, Di Loreto PC, O'Lane JM. A study of fetal heart rate acceleration patterns. *Obstet Gynecol* 1975;45:142-6.